Equine Preventative Care Health Plan

FOALS (< 1 year of age)

Immediate (within 12-24 hours) Newborn check-up:

- General physical exam (General health, heart defects, anatomical abnormalities, etc.)
- Pull blood to check for adequate antibody level (best if within 12 hours, 12-24 is adequate)
- Dip naval
- Vaccinate with tetanus antitoxin
- Enema
- Deworm Mare (recommendations)
- Check Mare (adequate milk supply, passage of placenta, body condition, etc.)

Additional Checkups at 3 and 6 months, to assess: Growth, Leg angles, etc.

Vaccinations:

3-4 months	4-6 months	6-9 months
EWT-WN* Influenza EHV/Rhinopneumonitis	EWT-WN* Influenza EHV/Rhinopneumonitis Rabies	EWT-WN* Influenza EHV/Rhinopneumonitis
	Recommended in some cases:**	Recommended in some cases:**
	Strangles	Strangles

^{*} Eastern and Western Encephalitis, Tetanus, Influenza virus

All equine vaccination recommendations conform to AAEP quidelines.

De-worming:

Every 2 months for first six months and again at 9 months de-worm with a mild paste medication such as strongid, panacur, or ivomec.

At one year of age perform <u>Equine Parasite Analysis</u>, with subsequent de-worming based on results.

^{**} See "Equine Vaccines" for detailed information.